

MODULI OF HOLOMORPHIC FUNCTIONS AND LOGARITHMICALLY CONVEX RADIAL WEIGHTS

EVGENY ABAKUMOV AND EVGUENI DOUBTSOV

ABSTRACT. Let $H(\mathbb{D})$ denote the space of holomorphic functions on the unit disk $\mathbb{D}$. We characterize those radial weights w on $\mathbb{D}$, for which there exist functions $f, g \in H(\mathbb{D})$ such that the sum $|f| + |g|$ is equivalent to w . Also, we obtain similar results in several complex variables for circular, strictly convex domains with smooth boundary.

1. INTRODUCTION

1.1. Growth spaces. Let $H(\mathbb{D})$ denote the space of holomorphic functions on the unit disk $\mathbb{D}$. Consider a weight function w , that is, a non-decreasing, continuous, unbounded function $w : [0, 1) \rightarrow (0, +\infty)$. We extend w to a radial weight on $\mathbb{D}$ setting $w(z) = w(|z|)$, $z \in \mathbb{D}$. By definition, the growth space $\mathcal{A}^w(\mathbb{D})$ consists of $f \in H(\mathbb{D})$ such that

$$(1.1) \quad |f(z)| \leq Cw(z), \quad z \in \mathbb{D},$$

for some constant $C > 0$.

Given functions $u, v : \mathbb{D} \rightarrow (0, +\infty)$, we write $u \asymp v$ and we say that u and v are equivalent if

$$C_1 u(z) \leq v(z) \leq C_2 u(z), \quad z \in \mathbb{D},$$

for some constants $C_1, C_2 > 0$. The definition of equivalent weight functions is analogous. If w_1 and w_2 are equivalent radial weights, then the identities

$$\|f\|_{\mathcal{A}^{w_j}(\mathbb{D})} = \sup_{z \in \mathbb{D}} \frac{|f(z)|}{w_j(z)}, \quad j = 1, 2,$$

define equivalent norms on the Banach space $\mathcal{A}^{w_1}(\mathbb{D}) = \mathcal{A}^{w_2}(\mathbb{D})$.

1.2. Motivations. In various applications, it is useful to have test functions $f \in \mathcal{A}^w(\mathbb{D})$ for which the reverse of estimate (1.1) holds, in a sense. The starting point for the present paper is the following reverse estimate obtained by W. Ramey and D. Ullrich [17] with the help of lacunary series:

Theorem 1.1 (cf. [17, Proposition 5.4]). *Let $w(t) = \frac{1}{1-t}$, $0 \leq t < 1$. There exist functions $f_1, f_2 \in \mathcal{A}^w(\mathbb{D})$ such that*

$$|f_1(z)| + |f_2(z)| \geq w(|z|)$$

for all $z \in \mathbb{D}$.

2010 *Mathematics Subject Classification.* Primary 30H99, 32A37; Secondary 42A55, 47B38.

Key words and phrases. Growth space, log-convex radial weight, circular domain.

The second author was partially supported by RFBR (grant No. 11-01-00526-a).

One may consider Theorem 1.1 as a particular solution of the following approximation problem:

Given a radial weight w on $\mathbb{D}$, find $f_1, f_2 \in H(\mathbb{D})$ such that

$$|f_1| + |f_2| \asymp w.$$

The above problem has been solved recently for various explicit radial weights. Clearly, the required property still holds if w is replaced by an equivalent radial weight. To the best knowledge of the authors, the largest class of weight functions $w(t)$ is considered in [1], where the direct analog of Theorem 1.1 is proved under assumption that $w(t)$ has the following doubling property:

$$(1.2) \quad w(1 - s/2) < Aw(1 - s), \quad 0 < s \leq 1,$$

for some constant $A > 1$. As in [17], the arguments in [1] use lacunary series; see [9] for a different proof. A similar result was independently obtained in [13]. See [1, 13] for further references.

Essentially, property (1.2) means that $w(t)$ grows sufficiently slowly as $t \rightarrow 1-$. So, it is natural to ask whether a growth restriction is crucial for the corresponding results. In the present paper, we characterize those w , for which the approximation problem under consideration is solvable. In particular, analogues of Theorem 1.1 hold for rapidly growing radial weights. However, there are natural restrictions on the regularity of w .

1.3. Main results. Let $w : [0, 1) \rightarrow (0, +\infty)$ be a weight function. Often, w is called log-convex if $\log w$ is convex. In this paper, we use a different definition related to Hadamard's three-circles theorem. Namely, w is said to be *log-convex* if $\log w(t)$ is a convex function of $\log t$, $0 < t < 1$. If $\log w(t)$ is a convex function of t , then w is log-convex, but the converse is not true; see subsection 2.2 for a discussion.

1.3.1. Unit disk.

Theorem 1.2. *Let w be a radial weight on $\mathbb{D}$. Then the following properties are equivalent:*

$$(1.3) \quad \text{there exist } f_1, f_2 \in H(\mathbb{D}) \text{ such that } |f_1| + |f_2| \asymp w;$$

$$(1.4) \quad w(t) \text{ is equivalent to a log-convex weight function on } [0, 1).$$

In other words, the above theorem characterizes those w for which an analog of Theorem 1.1 holds.

It is worth to mention the ideas behind the proof of Theorem 1.2. The implication (1.3) $\Rightarrow$ (1.4) is a corollary of Hadamard's three-circles theorem. To prove the reverse implication, we use a geometric construction that generates f_1 and f_2 as appropriate lacunary series. The construction is inspired by Borichev's work [5] on Fock-type spaces. Note that the series used in [1, 13, 17] are Hadamard lacunary. In contrast, for rapidly growing weight functions, we obtain weakly lacunary series: the corresponding frequencies grow slower than any geometric progression.

Remark 1.1. Condition (1.4) is not a growth restriction: given a weight function v , there exists a log-convex weight function w such that $w(t) \geq v(t)$, $0 \leq t < 1$. As mentioned above, (1.4) is a natural regularity condition.

Remark 1.2. Theorem 1.2 guarantees that property (1.4) also characterizes those radial weights which are equivalent to $\max(|f_1|, |f_2|)$ for some $f_1, f_2 \in H(\mathbb{D})$.

Remark 1.3. Given an integer M , one may consider the following problem: Characterize those radial weights w , for which there exist $f_1, \dots, f_M \in H(\mathbb{D})$ such that

$$|f_1| + \dots + |f_M| \asymp w.$$

The problem degenerates for $M = 1$: by the maximum modulus principle, no radial weight is equivalent to the modulus of a holomorphic function. For $M = 2, 3, \dots$, the solution of the problem does not depend on M and it is given by (1.4); see Remark 2.1.

1.3.2. Circular domains. We extend Theorem 1.2 to certain circular domains in $\mathbb{C}^d$, $d \in \mathbb{N}$. In particular, an analog of Theorem 1.2 holds for the unit ball of $\mathbb{C}^d$.

Assume that $\mathcal{D} \subset \mathbb{C}^d$ is a bounded, circular, strictly convex domain with $\mathcal{C}^2$ -boundary. Let $r_{\mathcal{D}}(z)$ denote the Minkowski functional on $\mathcal{D}$, that is,

$$r_{\mathcal{D}}(z) = \inf\{\rho > 0 : z \in \rho\mathcal{D}\}.$$

If $\mathcal{D}$ is the unit ball, then $r_{\mathcal{D}}(z) = |z|$.

Let $H(\mathcal{D})$ denote the space of holomorphic functions on $\mathcal{D}$. Given a weight function $w : [0, 1) \rightarrow (0, +\infty)$, the growth space $\mathcal{A}^w(\mathcal{D})$ consists of $f \in H(\mathcal{D})$ such that

$$\|f\|_{\mathcal{A}^w(\mathcal{D})} = \sup_{z \in \mathcal{D}} \frac{|f(z)|}{w(r_{\mathcal{D}}(z))} < \infty.$$

We obtain the following extension of Theorem 1.2 to several complex variables:

Theorem 1.3. *Let $\mathcal{D} \subset \mathbb{C}^d$ be a bounded, circular, strictly convex domain with $\mathcal{C}^2$ -boundary. If $w : [0, 1) \rightarrow (0, +\infty)$ is a log-convex weight function, then there exist functions $f_m \in \mathcal{A}^w(\mathcal{D})$, $1 \leq m \leq M = M(\mathcal{D})$ such that*

$$(1.5) \quad \sum_{m=1}^M |f_m(z)| \geq w(r_{\mathcal{D}}(z)), \quad z \in \mathcal{D}.$$

Conversely, if (1.5) holds for some $M \in \mathbb{N}$ and some $f_1, \dots, f_M \in H(\mathcal{D})$, then $w(t)$ is equivalent to a log-convex weight function.

Remark 1.4. Recall that, by Theorem 1.2 and Remark 1.3, the optimal value of M for $\mathbb{D}$ is equal to 2. It would be interesting to find the optimal value of $M(B_d)$ for the unit ball B_d of $\mathbb{C}^d$, $d \geq 2$.

1.4. Applications. Applications of Theorems 1.2 and 1.3 are mostly similar to those presented in [1] for the doubling weight functions: Carleson measures for $\mathcal{A}^w(\mathcal{D})$, composition and multiplication operators, Volterra type operators and extended Cesàro operators.

1.5. Organization of the paper. Section 2 is devoted to the proof of Theorem 1.2. Circular domains in $\mathbb{C}^d$ are considered in Section 3. Applications are discussed in the final Section 4.

Acknowledgement. The authors are grateful to Alexander Borichev for useful discussions and elucidating comments.

2. $\mathcal{D}$ IS THE UNIT DISK

In this section, we prove Theorem 1.2.

2.1. Equivalence to a log-convex weight function is necessary. Given a continuous function $u : \mathbb{D} \rightarrow (0, +\infty)$, put

$$M_u(r) = \max_{|z|=r} u(z), \quad 0 \leq r < 1.$$

Now, suppose that w is a radial weight, $f_1, f_2 \in H(\mathbb{D})$ and $|f_1| + |f_2| \asymp w$. Without loss of generality, we also assume that $f_j(0) \neq 0$ for $j = 1, 2$. By Hadamard's three circles theorem, $M_{|f_1|}$ and $M_{|f_2|}$ are log-convex functions on the interval $[0, 1]$; see, e.g., [6, Ch. 6, Sect. 3.13]. Hence, the sum $M_{|f_1|} + M_{|f_2|}$ is also log-convex (see, for example, [12, p. 51]).

Observe that

$$M_{|f_j|} \leq M_{|f_1|+|f_2|} \leq M_{|f_1|} + M_{|f_2|}, \quad j = 1, 2.$$

Thus, we have the following equivalences on $[0, 1]$:

$$w \asymp M_{|f_1|+|f_2|} \asymp M_{|f_1|} + M_{|f_2|}.$$

Since the function $M_{|f_1|} + M_{|f_2|}$ is log-convex, the implication (1.3) $\Rightarrow$ (1.4) is proved.

Remark 2.1. Let w be a radial weight such that

$$w \asymp |f_1| + \dots + |f_K|$$

for some $f_1, \dots, f_K \in H(\mathbb{D})$, $K \in \mathbb{N}$. Repeating the arguments used above for $K = 2$, we deduce that $w(t)$ is equivalent to a log-convex weight function.

The rest of the section is devoted to a constructive proof of the implication (1.4) $\Rightarrow$ (1.3).

2.2. Log-convex weight functions: preliminaries. Let $w : [0, 1] \rightarrow (0, +\infty)$ be a log-convex weight function. Recall that, by definition, $\log w(t)$ is a convex function of $\log t$, that is,

$$\Phi(x) = \Phi_w(x) = \log w(e^x), \quad x \in (-\infty, 0),$$

is a convex function.

Remark 2.2. It is natural to compare the above property and the following one: $\log w(t)$ is a convex function of t . The latter property implies that Φ_w is convex, as the composition of two increasing convex functions. The reverse implication does not hold. Moreover, there exist log-convex weight functions that are not even equivalent to the exponent of a convex function.

In what follows, we argue in terms of the function Φ . Observe that $w(t) = \exp(\Phi(\log t))$, $0 < t < 1$.

To prove the implication (1.4) $\Rightarrow$ (1.3), we may replace w by an equivalent weight function. So, without loss of generality, we assume that Φ is a strictly convex $\mathcal{C}^2$ -function. In particular, the tangent to the graph of Φ is unique at each point $(x, \Phi(x))$, $x < 0$. Also, below we repeatedly use the following property without explicit reference: the slope of the tangent to the graph of Φ at $(x, \Phi(x))$ is a strictly increasing function of $x \in (-\infty, 0)$.

Finally, note that $\log v(e^x) = \log a + \beta x$ for $v(t) = at^\beta$, $a > 0$, $\beta > 0$. This observation allows to reduce the proof of the implication (1.4) $\Rightarrow$ (1.3) to certain manipulations with linear functions.




FIGURE 1. Basic induction construction

2.3. Basic induction construction. Fix a number $x_0 \in (-\infty, 0)$ and a parameter $h > 0$. By induction, we construct linear functions $\ell_k(x)$ and numbers $x_k \in (x_0, 0)$, $k = 1, 2, \dots$, such that

- $\ell_k(x)$ is a tangent to the graph of $\Phi(x)$;
- $\ell_k(x)$ intersects the graph of $\Phi(x) - h$ at the points whose x -coordinates are x_{k-1} and x_k , $x_{k-1} < x_k$.

Note that $\ell_k(x)$ and x_k are uniquely defined by the above properties. So, the induction construction proceeds. See Figure 1.

Given $\ell_k(x)$, we define parameters $a_k > 0$ and $\beta_k > 0$ by the following identity:

$$\ell_k(x) = \log a_k + \beta_k x, \quad x \in \mathbb{R}.$$

Note that the sequences $\{x_k\}_{k=0}^\infty$ and $\{\beta_k\}_{k=1}^\infty$ monotonically increase, $x_k \rightarrow 0$ and $\beta_k \rightarrow \infty$ as $k \rightarrow \infty$. Also, we use the following brief notation: $t_k = \exp(x_k)$, $k = 0, 1, \dots$. Hence, the positive numbers t_k monotonically increase to 1 as $k \rightarrow \infty$.

Formally, the above construction works for any $h > 0$. In applications, we use a sufficiently large parameter h , say, $h = 2$, in the case of the unit disk.

2.4. Auxiliary estimates.

Lemma 2.1. *Let the numbers x_k , $k = 0, 1, \dots$, and the linear functions ℓ_k , $k = 1, 2, \dots$, be those introduced in subsection 2.3. Then*

$$(2.1) \quad \ell_{k+1}(x) \geq \ell_{k+2}(x) + h \quad \text{for all } x, \quad x_0 \leq x \leq x_k, \quad k \geq 0;$$

$$(2.2) \quad \ell_{k+1}(x) \geq \ell_k(x) + h \quad \text{for all } x, \quad x_{k+1} \leq x < 0, \quad k \geq 1.$$

Proof. We verify property (2.1). The proof of (2.2) is analogous, so we omit it.

Let $k = 1, 2, \dots$ and let x'_k denote the x -coordinate of the point at which ℓ_{k+1} is tangent to the graph of Φ . Remark that $x_k < x'_k < x_{k+1}$ by the definition of ℓ_{k+1} .

Now, consider three linear functions related to the convex function $\Phi - h$. First, $\ell_{k+1} - h$ is tangent to the graph of $\Phi - h$ at the point $(x'_k, (\Phi - h)(x'_k))$. Second, let L_{k+2} denote the tangent line to the graph of $\Phi - h$ at $(x_{k+1}, (\Phi - h)(x_{k+1}))$. Third, recall that ℓ_{k+2} intersects the graph of $\Phi - h$ at the points $(x_{k+1}, (\Phi - h)(x_{k+1}))$ and $(x_{k+2}, (\Phi - h)(x_{k+2}))$.

We have

$$(2.3) \quad (\ell_{k+1} - h)(x'_k) = (\Phi - h)(x'_k) > L_{k+2}(x'_k),$$

because $x'_k < x_{k+1}$ and $\Phi - h$ is convex.

Next, $L_{k+2}(x_{k+1}) = \ell_{k+2}(x_{k+1})$ and the slope of the linear function L_{k+2} is smaller than that of ℓ_{k+2} by the definitions of L_{k+2} and ℓ_{k+2} . Hence, the inequality $x'_k < x_{k+1}$ guarantees that

$$(2.4) \quad L_{k+2}(x'_k) > \ell_{k+2}(x'_k).$$

By (2.3) and (2.4), we obtain $(\ell_{k+1} - h)(x'_k) > \ell_{k+2}(x'_k)$. Therefore,

$$(\ell_{k+1} - h)(x) > \ell_{k+2}(x) \quad \text{for all } x \leq x'_k,$$

because the slope of $\ell_{k+1} - h$ is smaller than that of ℓ_{k+2} . Recall that $x_k < x'_k$, thus, the lemma is proved. $\square$

Lemma 2.2. *Let the numbers x_k , t_k , $k = 0, 1, \dots$, and the linear functions ℓ_k , $k = 1, 2, \dots$, be those introduced in subsection 2.3. Assume that $h \geq 2$. Then, for $k = 1, 2, \dots$,*

- (i) $a_k t^{\beta_k} \leq w(t)$, $t \in [t_0, 1]$;
- (ii) $e^{-h} w(t) \leq a_k t^{\beta_k}$, $t \in [t_{k-1}, t_k]$;
- (iii) $\sum_{m \geq 1, |m-k| \geq 2} a_m t^{\beta_m} < \frac{1}{2} a_k t^{\beta_k}$, $t \in [t_{k-1}, t_k]$.

Proof. Let $k \in \mathbb{N}$. We have $\ell_k(x) \leq \Phi(x)$ for $x_0 \leq x < 0$. Hence, taking the exponentials, we obtain (i) by the definitions of Φ , ℓ_k , a_k and β_k . Analogously, the inequality $\Phi(x) - h \leq \ell_k(x)$, $x \in [x_{k-1}, x_k]$, implies (ii).

It remains to prove (iii). First, fix a $k \geq 1$ and assume that $m \geq k + 2$. For $x_0 \leq x \leq x_k$, we have

$$\ell_k(x) - \ell_m(x) \geq \ell_{k+1}(x) - \ell_m(x) = \sum_{j=k+1}^{m-1} [\ell_j(x) - \ell_{j+1}(x)].$$

Consider the latter sum. For $k+1 \leq j \leq m-1$, property (2.1) guarantees that $\ell_j(x) - \ell_{j+1}(x) \geq h$ for all $x_0 \leq x \leq x_{j-1}$, hence, for all $x_0 \leq x \leq x_k$. In sum, we have

$$\ell_k(x) - \ell_m(x) \geq h(m - k - 1) \quad \text{for } x_0 \leq x \leq x_k.$$

Taking the exponentials and using the definitions of ℓ_m and ℓ_k , we obtain

$$a_m t^{\beta_m} \leq a_k t^{\beta_k} \frac{1}{e^{h(m-k-1)}} \quad \text{for } t_0 \leq t \leq t_k.$$

Thus,

$$(2.5) \quad \sum_{m \geq k+2} a_m t^{\beta_m} \leq a_k t^{\beta_k} \left(\frac{1}{e^h} + \frac{1}{e^{2h}} + \frac{1}{e^{3h}} + \dots \right) \quad \text{for } t_0 \leq t \leq t_k.$$

Second, fix $k \geq 3$ and assume that $1 \leq m \leq k-2$. Essentially, we argue as above, replacing (2.1) by (2.2). Namely, for $0 > x \geq x_{k-1}$, we have

$$\ell_k(x) - \ell_m(x) \geq \ell_{k-1}(x) - \ell_m(x) = \sum_{j=m}^{k-2} [\ell_{j+1}(x) - \ell_j(x)].$$

For $m \leq j \leq k-2$, property (2.2) guarantees that $\ell_{j+1}(x) - \ell_j(x) \geq h$ for all $0 > x \geq x_{k-1}$. In sum, we have

$$\ell_m(x) \leq \ell_k(x) - h(k-m-1) \quad \text{for } 0 > x \geq x_{k-1}.$$

Taking the exponentials and summing over $m = 1, \dots, k-2$, we obtain

$$(2.6) \quad \sum_{m=1}^{k-2} a_m t^{\beta_m} \leq a_k t^{\beta_k} \left(\frac{1}{e^h} + \frac{1}{e^{2h}} + \frac{1}{e^{3h}} + \dots \right) \quad \text{for } 1 > t \geq t_{k-1}.$$

In sum, (iii) holds by (2.5) and (2.6), because $h \geq 2$. The proof of the lemma is complete. $\square$

2.5. Proof of (1.4) $\Rightarrow$ (1.3). Fix a parameter $h \geq 2$ and a number $x_0 \in (-\infty, 0)$ such that $t_0 = \exp(x_0) > \frac{9}{10}$. The basic induction construction provides numbers x_k, t_k, β_k and $a_k, k = 1, 2, \dots$. Put $n_k = [\beta_k] + 1, k = 1, 2, \dots$, where $[\beta_k]$ denote the integer part of β_k . Without loss of generality, we assume that x_0 is so close to 0 that $n_k < n_{k+1}$ for all $k = 1, 2, \dots$.

Put

$$g_1(z) = \sum_{j=0}^{\infty} a_{2j+1} z^{n_{2j+1}};$$

$$g_2(z) = \sum_{j=1}^{\infty} a_{2j} z^{n_{2j}}.$$

The estimates below guarantee that g_1 and g_2 are well-defined functions of $z \in \mathbb{D}$. In fact, we claim that

$$(2.7) \quad \frac{2}{5} e^{-h} w(z) < |g_1(z)| + |g_2(z)| < 4w(z), \quad t_0 < |z| < 1.$$

To prove the above estimates, put $t = |z|$. For $t \in (\frac{9}{10}, 1)$, we have

$$1 < \frac{a_k t^{\beta_k}}{a_k t^{n_k}} \leq \frac{1}{t} \leq \frac{10}{9}.$$

Therefore, properties (i–iii) from Lemma 2.2 imply the following inequalities:

$$\begin{aligned} \text{(i')} \quad & a_k t^{n_k} \leq w(t), \quad t \in [t_0, 1]; \\ \text{(ii')} \quad & \frac{9}{10} e^{-h} w(t) \leq a_k t^{n_k}, \quad t \in [t_{k-1}, t_k]; \\ \text{(iii')} \quad & \end{aligned}$$

$$\sum_{m \geq 1, |m-k| \geq 2} a_m t^{n_m} < \sum_{m \geq 1, |m-k| \geq 2} a_m t^{\beta_m} < \frac{1}{2} a_k t^{\beta_k} \leq \frac{5}{9} a_k t^{n_k}, \quad t \in [t_{k-1}, t_k].$$

Now, assume that $t \in [t_{k-1}, t_k]$ for some odd number k . On the one hand,

$$\begin{aligned}
\frac{2}{5}e^{-h}w(t) &\stackrel{(ii')}{<} \frac{4}{9}a_k t^{n_k} \\
&\stackrel{(iii')}{<} a_k t^{n_k} - \sum_{m \geq 1, |m-k| \geq 2} a_m t^{n_m} \\
&\leq \left| \sum_{j=0}^{\infty} a_{2j+1} z^{n_{2j+1}} \right| \\
&= |g_1(z)| \\
&\leq |g_1(z)| + |g_2(z)|.
\end{aligned}$$

On the other hand, if $k \geq 3$, then

$$\begin{aligned}
|g_1(z)| + |g_2(z)| &\leq \sum_{m \geq 1} a_m t^{n_m} \\
&= \left(a_k t^{n_k} + \sum_{m \geq 1, |m-k| \geq 2} a_m t^{n_m} \right) + (a_{k-1} t^{n_{k-1}} + a_{k+1} t^{n_{k+1}}) \\
&\stackrel{(i'), (iii')}{\leq} \left(1 + \frac{5}{9} \right) a_k t^{n_k} + 2w(t) \\
&\stackrel{(i')}{<} 4w(t).
\end{aligned}$$

For $k = 1$, the above arguments are even more simple.

If k is even, then we have analogous estimates. So, (2.7) holds.

To finish the proof, we modify the functions g_1 and g_2 . Namely, we have $g_1(z) = z^{n_1} G_1(z)$, where $G_1 \in H(\mathbb{D})$ and $G_1(0) \neq 0$. Clearly, $G_1 \in \mathcal{A}^w(\mathbb{D})$ and $|G_1(z)| \geq |g_1(z)|$ for all $z \in \mathbb{D}$. Therefore,

$$\frac{2}{5}e^{-h}w(z) < |G_1(e^{i\theta}z)| + |g_2(z)|, \quad t_0 < |z| < 1,$$

for all $\theta \in [0, 2\pi)$. The functions G_1 and g_2 have a finite number of zeros in the disk $\{|z| \leq t_0\}$ and $G_1(0) \neq 0$; hence, the required property

$$|f_1(z)| + |f_2(z)| \asymp w(z), \quad z \in \mathbb{D},$$

holds for $f_2(z) = g_2(z)$ and $f_1(z) = G_1(e^{i\theta}z)$ with an appropriate $\theta \in [0, 2\pi)$. So, (1.4) implies (1.3). The proof of Theorem 1.2 is finished.

3. $\mathcal{D}$ IS A CIRCULAR DOMAIN

3.1. Equivalence to a log-convex weight function. This question reduces to the corresponding implication in the unit disk. Indeed, assume that (1.5) holds. Fix a point $\zeta \in \partial\mathcal{D}$. For $m = 1, \dots, M$, consider the slice-functions $g_m(\lambda) = f_m(\lambda\zeta)$, $\lambda \in \mathbb{D}$. We have $g_m \in H(\mathbb{D})$, $m = 1, \dots, M$, and

$$|g_1(\lambda)| + \dots + |g_M(\lambda)| \asymp w(r_{\mathcal{D}}(\lambda\zeta)) = w(|\lambda|), \quad \lambda \in \mathbb{D}.$$

Hence, as indicated in Remark 2.1, $w(t)$ is equivalent to a log-convex weight function.

3.2. Aleksandrov–Ryll–Wojtaszczyk polynomials. The proof of the existence part in Theorem 1.3 will be based on lacunary series of special homogeneous holomorphic polynomials (cf. [2, 18]).

Theorem 3.1. *Given a domain $\mathcal{D} \subset \mathbb{C}^d$, there exist $\delta = \delta(\mathcal{D}) \in (0, 1)$ and $Q = Q(\mathcal{D}) \in \mathbb{N}$ with the following properties: for every $n \in \mathbb{N}$, there exist homogeneous holomorphic polynomials $W_q[n]$ of degree n , $1 \leq q \leq Q$, such that*

$$(3.1) \quad \|W_q[n]\|_{L^\infty(\partial\mathcal{D})} \leq 1;$$

$$(3.2) \quad \max_{1 \leq q \leq Q} |W_q[n](\zeta)| \geq \delta \quad \text{for all } \zeta \in \partial\mathcal{D}.$$

As observed in [8], to prove Theorem 3.1, it suffices to repeat *mutatis mutandis* the arguments used in [11, Theorem 2.6].

3.3. Auxiliary estimates. We need a modification of property (iii) from Lemma 2.2.

Lemma 3.2. *Let $\delta \in (0, 1)$. Then there exists $h(\delta) \geq 2$ with the following property: Let the numbers $x_k, t_k, k = 0, 1, \dots$, and the linear functions $\ell_k, k = 1, 2, \dots$, be those introduced in subsection 2.3. Assume that $h \geq h(\delta)$. Then*

$$(iii_\delta) \quad \sum_{m \geq 1, |m-k| \geq 2} a_m t^{\beta_m} < \frac{\delta}{2} a_k t^{\beta_k}, \quad t \in [t_{k-1}, t_k],$$

for $k = 1, 2, \dots$.

Proof. The argument coincides with that used in the proof of Lemma 2.2. Indeed, if $h \geq 2$ is sufficiently large, then properties (2.5) and (2.6) imply (iii_δ). $\square$

3.4. Proof of Theorem 1.3: existence part. Let $w : [0, 1) \rightarrow (0, +\infty)$ be a log-convex weight function and let $\Phi(x) = \log w(e^x)$, $x \in (-\infty, 0)$.

We modify the arguments used in the case of the unit disk, replacing the monomials z^n , $n = 1, 2, \dots$, by the Aleksandrov–Ryll–Wojtaszczyk polynomials $W_q[n]$, $q = 1, \dots, Q$, $n = 1, 2, \dots$.

So, let the constant $\delta \in (0, 1)$ be that provided by Theorem 3.1 and let the constant $h(\delta) \geq 2$ be that provided by Lemma 3.2. Fix a parameter $h \geq h(\delta)$ and a number $x_0 \in (-\infty, 0)$ such that $t_0 = \exp(x_0) > \frac{9}{10}$. The basic induction construction provides numbers x_k, t_k, β_k and $a_k, k = 1, 2, \dots$.

For $s = 0, 1$, put

$$f_{q+sQ}(z) = \sum_{j=0}^{\infty} a_{2j+1+s} W_q[n_{2j+1+s}](z), \quad z \in \mathcal{D}, \quad q = 1, \dots, Q.$$

where $n_k = [\beta_k] + 1, k = 1, 2, \dots$.

We adapt the argument used to prove (2.7). So, let $t = r_{\mathcal{D}}(z)$. Properties (i–ii) from Lemma 2.2 and property (iii_δ) from Lemma 3.2 imply the following inequalities:

$$\begin{aligned} (i') \quad & a_k t^{n_k} \leq w(t), \quad t \in [t_0, 1); \\ (ii') \quad & \frac{9}{10} e^{-h} w(t) \leq a_k t^{n_k}, \quad t \in [t_{k-1}, t_k]; \\ (iii'_\delta) \quad & \end{aligned}$$

$$\sum_{m \geq 1, |m-k| \geq 2} a_m t^{n_m} < \sum_{m \geq 1, |m-k| \geq 2} a_m t^{\beta_m} < \frac{\delta}{2} a_k t^{\beta_k} \leq \frac{5\delta}{9} a_k t^{n_k}, \quad t \in [t_{k-1}, t_k].$$

First, remark that (iii_δ') is more stringent than (iii'). So, f_{q+sQ} are well-defined functions of $z \in \mathcal{D}$. Moreover, combining estimate (3.1) and the arguments from the proof of Theorem 1.2, we obtain $f_{q+sQ} \in \mathcal{A}^w(\mathcal{D})$, $s = 0, 1$, $q = 1, \dots, Q$.

Second, we claim that

$$(3.3) \quad \frac{2}{5}e^{-h}w(t) < \sum_{q=1}^Q |f_q(z)| + \sum_{q=1}^Q |f_{q+Q}(z)|, \quad t_0 < t < 1.$$

To prove the above estimate, assume that $t \in [t_{k-1}, t_k]$ for some odd number k . We have $z = t\zeta$ for some $\zeta \in \partial\mathcal{D}$. Applying (3.2), select $q(\zeta)$, $1 \leq q(\zeta) \leq Q$ such that

$$(3.4) \quad |W_{q(\zeta)}[n_k](\zeta)| \geq \delta.$$

Recall that $W_{q(\zeta)}[n]$ is a homogeneous polynomial of degree n . Thus, we obtain

$$\begin{aligned} \frac{2\delta}{5}e^{-h}w(t) &\stackrel{(ii')}{<} \frac{4\delta}{9}a_k t^{n_k} \\ &\stackrel{(iii_\delta')}{<} \delta a_k t^{n_k} - \sum_{m \geq 1, |m-k| \geq 2} a_m t^{n_m} \\ &\stackrel{(3.1) \text{ and } (3.4)}{\leq} \left| \sum_{j=0}^{\infty} a_{2j+1} W_{q(\zeta)}[n_{2j+1}](t\zeta) \right| \\ &= |f_{q(\zeta)}(t\zeta)| \\ &\leq \sum_{q=1}^Q |f_q(t\zeta)| + \sum_{q=1}^Q |f_{q+Q}(t\zeta)|. \end{aligned}$$

If k is even, then we have an analogous estimate. So, (3.3) holds. Putting $f_{2Q+1} \equiv 1$, we obtain the required estimate

$$w(r_{\mathcal{D}}(z)) \leq C \sum_{m=1}^{2Q+1} |f_m(z)|$$

for all $z \in \mathcal{D}$. The proof of Theorem 1.3 is finished.

4. APPLICATIONS

Theorem 1.3 allows to recover the applications presented in [1, Sections 3–7] for the doubling weight functions w . In this section, we consider several direct applications of Theorem 1.3, assuming that $w : [0, 1) \rightarrow (0, +\infty)$ is a log-convex weight function, unless otherwise stated.

4.1. Composition operators and Carleson measures. Suppose that $\mathcal{D} \subset \mathbb{C}^d$ is a bounded, circular, strictly convex domain with $\mathcal{C}^2$ -boundary.

4.1.1. Weighted composition operators. Given $g \in H(\mathcal{D})$ and a holomorphic mapping $\varphi : \mathcal{D} \rightarrow \mathcal{D}$, the weighted composition operator $C_\varphi^g : H(\mathcal{D}) \rightarrow H(\mathcal{D})$ is defined by the formula

$$(C_\varphi^g f)(z) = g(z)f(\varphi(z)), \quad f \in H(\mathcal{D}), \quad z \in \mathcal{D}.$$

If $g \equiv 1$, then C_φ^g is denoted by C_φ and it is called a composition operator. Various properties of C_φ are presented in monographs [7, 19].

Consider a linear space $\mathcal{Y}(\mathcal{D})$ which consists of functions $f : \mathcal{D} \rightarrow \mathbb{C}$. By definition, $\mathcal{Y}(\mathcal{D})$ is a lattice if the following implication holds:

Assume that $F \in \mathcal{Y}(\mathcal{D})$, $f : \mathcal{D} \rightarrow \mathbb{C}$ is a continuous function, and $|f(z)| \leq |F(z)|$ for all $z \in \mathcal{D}$. Then $f \in \mathcal{Y}(\mathcal{D})$.

Corollary 4.1. *Suppose that $g \in H(\mathcal{D})$, $\varphi : \mathcal{D} \rightarrow \mathcal{D}$ is a holomorphic mapping and $\mathcal{Y}(\mathcal{D})$ is a lattice. Then the weighted composition operator C_φ^g maps $\mathcal{A}^w(\mathcal{D})$ into $\mathcal{Y}(\mathcal{D})$ if and only if*

$$(4.1) \quad |g(z)|w(r_{\mathcal{D}}(\varphi(z))) \in \mathcal{Y}(\mathcal{D}).$$

Proof. Assume that C_φ^g maps $\mathcal{A}^w(\mathcal{D})$ to $\mathcal{Y}(\mathcal{D})$. Let the functions $f_m \in \mathcal{A}^w(\mathcal{D})$, $1 \leq m \leq M$, be those provided by Theorem 1.3. Applying inequality (1.5), we have

$$|g(z)|w(r_{\mathcal{D}}(\varphi(z))) \leq \sum_{m=1}^M |g(z)||f_m(\varphi(z))| = \sum_{m=1}^M |(C_\varphi^g f_m)(z)|, \quad z \in \mathcal{D}.$$

Since $\mathcal{Y}(\mathcal{D})$ is a lattice, we obtain (4.1). To prove the converse implication, suppose that (4.1) holds. If $f \in \mathcal{A}^w(\mathcal{D})$, then

$$|(C_\varphi^g f)(z)| \leq \|f\|_{\mathcal{A}^w(\mathcal{D})} |g(z)|w(r_{\mathcal{D}}(\varphi(z))) \in \mathcal{Y}(\mathcal{D}).$$

Hence, $C_\varphi^g f \in \mathcal{Y}(\mathcal{D})$ by the definition of a lattice. $\square$

4.1.2. Compact operators and Carleson measures. Given a lattice $\mathcal{Y}(\mathcal{D})$, characterizations of the compact operators $C_\varphi^g : \mathcal{A}^w(\mathcal{D}) \rightarrow \mathcal{Y}(\mathcal{D})$ depend on specific properties of $\mathcal{Y}(\mathcal{D})$ (see [1] for details). As an example, consider the case $\mathcal{Y}(\mathcal{D}) = L^p(\mathcal{D}, \mu)$.

Given a space $X \subset H(\mathcal{D})$ and $0 < p < \infty$, recall that a positive Borel measure μ on $\mathcal{D}$ is said to be p -Carleson for X if

the identity operator $I : X \rightarrow L^p(\mathcal{D}, \mu)$ is bounded.

Respectively, μ is called a vanishing p -Carleson measure for X if

the identity operator $I : X \rightarrow L^p(\mathcal{D}, \mu)$ is compact.

Corollary 4.2. *Suppose that $0 < p < \infty$ and μ is a positive Borel measure on $\mathcal{D}$. Then the following properties are equivalent:*

$$(4.2) \quad \mu \text{ is a } p\text{-Carleson measure for } \mathcal{A}^w(\mathcal{D});$$

$$(4.3) \quad \mu \text{ is a vanishing } p\text{-Carleson measure for } \mathcal{A}^w(\mathcal{D});$$

$$(4.4) \quad \int_{\mathcal{D}} w^p(r_{\mathcal{D}}(z)) d\mu(z) < \infty.$$

Proof. Properties (4.2) and (4.4) are equivalent by Corollary 4.1 with $\mathcal{Y}(\mathcal{D}) = L^p(\mathcal{D}, \mu)$, $g \equiv 1$ and $\varphi(z) \equiv z$. Also, standard arguments guarantee that (4.4) implies (4.3). Since (4.3) implies (4.2), the proof is finished. $\square$

4.2. Integral operators. Assume that $\mathcal{D}$ is the unit ball B_d of $\mathbb{C}^d$. Given $g \in H(B_d)$ and a holomorphic mapping $\varphi : B_d \rightarrow B_d$, the Volterra type operator $V_\varphi^g : H(B_d) \rightarrow H(B_d)$ is defined as

$$(V_\varphi^g f)(z) = \int_0^1 f(\varphi(tz)) \frac{\mathcal{R}g(tz)}{t} dt, \quad f \in H(B_d), \quad z \in B_d,$$

where

$$\mathcal{R}g(z) = \sum_{j=1}^d z_j \frac{\partial g}{\partial z_j}(z), \quad z \in B_d,$$

is the radial derivative of g . If $\varphi(z) \equiv z$, then V_φ^g is denoted by J_g and it is called an extended Cesàro operator (see [10]). In fact, if $n = 1$, then we have

$$(J_g f)(z) = \int_0^z f(w) g'(w) dw, \quad f \in H(B_1), \quad z \in B_1.$$

The above operator was introduced by Pommerenke [16] as a natural generalization of the classical Cesàro operator. For $n = 1$, various properties of the operator J_g are discussed in surveys [3, 20].

Direct calculations show that

$$(4.5) \quad \mathcal{R}V_\varphi^g f(z) = f(\varphi(z)) \mathcal{R}g(z), \quad z \in B_d,$$

for all $f, g \in H(B_d)$ (cf. [10]). Applying (4.5) and Corollary 4.1, we obtain the following fact.

Corollary 4.3. *Let $\mathcal{Y}(B_d)$ be a lattice on B_d . Then the operator $\mathcal{R}V_\varphi^g$ maps $\mathcal{A}^w(B_d)$ into $\mathcal{Y}(B_d)$ if and only if $|\mathcal{R}g(z)|w(|\varphi(z)|) \in \mathcal{Y}(B_d)$.*

Given a space $X \subset H(B_d)$, a typical problem is to characterize $g \in H(B_d)$ such that J_g is a bounded operator on X . Often the characterizing property is the following one:

$$\sup_{z \in B_d} |\mathcal{R}g(z)|(1 - |z|) < \infty,$$

that is, g is in the Bloch space $\mathcal{B}(B_d)$ (see, e.g., [4, 10]). Hence, it is interesting to find those X for which the answer is different. To give such examples, consider the unit disk $\mathbb{D} = B_1$ and the following exponential weight functions:

$$w_\alpha(r) = \exp\left(\frac{1}{(1-r)^\alpha}\right), \quad 0 \leq r < 1, \quad \alpha > 0.$$

Note that the above weight functions are not doubling.

Corollary 4.4. *Let $\alpha > 0$ and let $g \in H(\mathbb{D})$. The operator $J_g : \mathcal{A}^{w_\alpha}(\mathbb{D}) \rightarrow \mathcal{A}^{w_\alpha}(\mathbb{D})$ is bounded if and only if*

$$(4.6) \quad \sup_{z \in \mathbb{D}} |g(z)|(1 - |z|)^\alpha < \infty.$$

Proof. Let $f \in H(\mathbb{D})$. Theorem D from [15] guarantees that

$$J_g f \in \mathcal{A}^{w_\alpha}(\mathbb{D}) \Leftrightarrow \mathcal{R}J_g f \in \mathcal{A}^{w'_\alpha}(\mathbb{D}).$$

By Corollary 4.3, the latter property holds if and only if

$$\sup_{z \in \mathbb{D}} |\mathcal{R}g(z)| \frac{w_\alpha(|z|)}{w'_\alpha(|z|)} < \infty,$$

that is,

$$\sup_{z \in \mathbb{D}} |\mathcal{R}g(z)|(1 - |z|)^{\alpha+1} < \infty.$$

It remains to remark that the above property is equivalent to (4.6). $\square$

By definition, the weighted Bergman space $A_{w_\alpha}^p(\mathbb{D})$, $0 < p < \infty$, consists of $f \in H(\mathbb{D})$ such that

$$\int_{\mathbb{D}} |f(z)|^p \frac{d\nu(z)}{w_\alpha(|z|)} < \infty,$$

where ν denotes Lebesgue measure on $\mathbb{C}$.

Formally, one may consider $\mathcal{A}^{w_\alpha}(\mathbb{D})$ as the weighted Bergman space $A_{w_\alpha}^p(\mathbb{D})$ with $p = \infty$. So, given $\alpha > 0$, remark that property (4.6) also characterizes those $g \in H(\mathbb{D})$ for which the operator $J_g : A_{w_\alpha}^p(\mathbb{D}) \rightarrow A_{w_\alpha}^p(\mathbb{D})$, $0 < p < \infty$, is bounded (see [14]).

4.3. Other applications. If $\mathcal{D}$ is the unit disk $\mathbb{D}$, then Theorem 1.3 is also applicable to the weighted Bloch spaces; cf. [1, 13].

REFERENCES

1. E. Abakumov and E. Doubtsov, *Reverse estimates in growth spaces*, Math. Z. **271** (2012), no. 1-2, 399–413.
2. A. B. Aleksandrov, *Proper holomorphic mappings from the ball to the polydisk*, Dokl. Akad. Nauk SSSR **286** (1986), no. 1, 11–15 (Russian); English transl.: Soviet Math. Dokl. **33** (1) (1986), 1–5.
3. A. Aleman, *A class of integral operators on spaces of analytic functions*, Topics in complex analysis and operator theory, Univ. Málaga, Málaga, 2007, pp. 3–30.
4. A. Aleman and A. G. Siskakis, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. **46** (1997), no. 2, 337–356.
5. A. Borichev, *The polynomial approximation property in Fock-type spaces*, Math. Scand. **82** (1998), no. 2, 256–264.
6. J. B. Conway, *Functions of one complex variable*, second ed., Graduate Texts in Mathematics, vol. 11, Springer-Verlag, New York, 1978.
7. C. C. Cowen and B. D. MacCluer, *Composition operators on spaces of analytic functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1995.
8. E. Doubtsov, *Growth spaces on circular domains: composition operators and Carleson measures*, C. R. Math. Acad. Sci. Paris **347** (2009), no. 11-12, 609–612.
9. J. Gröhn, J. Á. Peláez, and J. Rättyä, *Jointly maximal products in weighted growth spaces*, preprint, arXiv:1210.3318.
10. Z. Hu, *Extended Cesàro operators on mixed norm spaces*, Proc. Amer. Math. Soc. **131** (2003), no. 7, 2171–2179.
11. P. Kot, *Homogeneous polynomials on strictly convex domains*, Proc. Amer. Math. Soc. **135** (2007), no. 12, 3895–3903.
12. S. G. Kreĭn, Yu. Ī. Petunĭn, and E. M. Semĕnov, *Interpolation of linear operators*, Translations of Mathematical Monographs, vol. 54, American Mathematical Society, Providence, R.I., 1982.
13. E. G. Kwon and M. Pavlović, *BiBloch mappings and composition operators from Bloch type spaces to BMOA*, J. Math. Anal. Appl. **382** (2011), no. 1, 303–313.
14. J. Pau and J. Á. Peláez, *Embedding theorems and integration operators on Bergman spaces with rapidly decreasing weights*, J. Funct. Anal. **259** (2010), no. 10, 2727–2756.
15. M. Pavlović and J. Á. Peláez, *An equivalence for weighted integrals of an analytic function and its derivative*, Math. Nachr. **281** (2008), no. 11, 1612–1623.
16. Ch. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, Comment. Math. Helv. **52** (1977), no. 4, 591–602.
17. W. Ramey and D. Ullrich, *Bounded mean oscillation of Bloch pull-backs*, Math. Ann. **291** (1991), no. 4, 591–606.

18. J. Ryll and P. Wojtaszczyk, *On homogeneous polynomials on a complex ball*, Trans. Amer. Math. Soc. **276** (1983), no. 1, 107–116.
19. J. H. Shapiro, *Composition operators and classical function theory*, Universitext: Tracts in Mathematics, Springer-Verlag, New York, 1993.
20. A. G. Siskakis, *Volterra operators on spaces of analytic functions—a survey*, Proceedings of the First Advanced Course in Operator Theory and Complex Analysis, Univ. Sevilla Secr. Publ., Seville, 2006, pp. 51–68.

UNIVERSITÉ PARIS-EST, LAMA (UMR 8050), UPEMLV, UPEC, CNRS, F-77454, MARNE-LA-VALLÉE, FRANCE

E-mail address: `Evgueni.Abakoumov@univ-mlv.fr`

ST. PETERSBURG DEPARTMENT OF V.A. STEKLOV MATHEMATICAL INSTITUTE, FONTANKA 27, ST. PETERSBURG 191023, RUSSIA

E-mail address: `dubtsov@pdmi.ras.ru`